



Comparative Study of Mineral and Phytochemical Analysis of Soil and *Lactuca sativa* Grown in the Vicinity of Cement Company of Northern Nigeria (Sokoto Cement) and Usmanu Danfodiyo University Sokoto (Kwankwalawa)

M. H. Dalhat^{1*}, A. R. Amale¹, M. Maimuna², I. Bashiru³ and K. Sirajo³

¹*Department of Biochemistry, Usmanu Danfodiyo University Sokoto, Nigeria.*

²*Department of Biological Sciences, Usmanu Danfodiyo University Sokoto, Nigeria.*

³*Department of Biochemistry, Sokoto State University, Nigeria.*

Authors' contributions

This work was carried out in collaboration between all authors. Author MHD designed the study. Authors MHD, ARA, MM and IB performed the statistical analysis. Author MHD wrote the protocol. Author ARA wrote the first draft of the manuscript. Authors MM, KS and IB managed the analyses of the study. Authors MHD, ARA and KS managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRB/2017/111329

Editor(s):

(1) Mohamed Fawzy Ramadan Hassanien, Professor, Biochemistry Department, Faculty of Agriculture, Zagazig University, Zagazig, Egypt.

Reviewers:

(1) Phyu Phyu Myint, University of Yangon, Myanmar.

(2) Ningappa M. Rolli, BLDEA,s Degree College Jamkhandi, India.

(3) Poliana Guerino Marson, Federal University of Tocantins, Brazil.

Complete Peer review History: <http://www.sciencedomain.org/review-history/22485>

Original Research Article

Received 7th December 2017
Accepted 23rd December 2017
Published 28th December 2017

ABSTRACT

Environmental pollution is a major issue which confronts industry and business in today's world on daily basis. Industrial activities are the leading cause of metals emission, often associated with soil and plant metal concentration in adjacent regions. Cement industry is one of the 17 most polluting industries listed by the central pollution control board (CPCB). Impact of dust deposition from Cement Company of Northern Nigeria on the proximate and phytochemical concentrations of lettuce (*Lactuca sativa*) was studied. A comparative study of heavy metal concentration and phytochemicals of *Lactuca sativa* and soil samples from Kalambaina (Industrial area) and

*Corresponding author: E-mail: mahmoodalhat@gmail.com;

Kwalkwalawa (non-Industrial area) were estimated using atomic absorption spectrophotometry (AAS) and standard analytical procedures respectively. Result of quantitative phytochemical analysis revealed significant difference ($P < 0.05$) in all parameters. Heavy metal values of Pb(0.012 ± 0.002 mg/g), Zn(0.043 ± 0.003 mg/g), and Ca(706.860 ± 14.980 mg/g) in *Lactuca sativa* collected from Kalambaina revealed significant difference ($P < 0.05$) when compare to samples collected from Kwalkwalawa and WHO standard. In addition, the heavy metal concentration in soil collected from Kalambaina showed significant difference ($P < 0.05$) when compare to samples collected from Kwalkwalawa; with the highest value recorded in Ca (974.25 ± 48 mg/g) which might be as a result of activities in the cement industry. Conclusively, plants grown at cement industries might not be safe for consumption.

Keywords: Environmental pollution; micronutrients; phytochemicals; atomic absorption spectrophotometry; ash; moisture.

1. INTRODUCTION

Environmental pollution is a major issue associated with industry and business in today's world on daily basis. Various environmental components such as water, air, soil, and vegetation are degraded as a result of Industrial and human activities [1]. The environmental pollution as a result of cement industry could be defined as the adverse effect induced on water, air and land through various activities, starting from mining activity of the raw material (lime stone, dolomite etc.) up to its crushing, grinding, and other processes developing in a cement industry [2].

Cement is a fine, gray or white powder which is largely made up of Cement Kiln Dust (CKD), a by-product of the final cement product, usually stored as wastes in open-pits and landfills. Exposure to cement dust for a short period may not cause serious problem, however prolonged exposure can cause serious irreversible damage to plants and animals. Dust which is emitted during cement processes are eventually deposited on soil, sediment, water and plants. The dust emissions from cement and other related industries therefore have to be given attention for control [3]. This is necessary in view of the pollution load and its impact on the environment. Pollutants have devastating effects on plants when taken by direct absorption through leaves or by water through the roots [4]. Heavy metals are natural constituents in nature, usually occurring in low concentration under normal conditions. Heavy metals contamination of vegetables cannot be underestimated as these food stuffs are important components of human diet [5]. Excessive concentration of heavy metals; such as Hg, Cu, Pb, Cd and Ni from pollutants may result in phytotoxicity in plants [5].

Plants are nature's gift to humans in terms of providing us with food, oxygen, as well as shelter. Since time immemorial [6], they have served as the first line of defense used by our forefathers to fight diseases such as Diarrhea, Cholera and Malaria [1]. Plants are conveniently separated into those which are edible, those which serve as a source of medicine or spices, and those that are of ornamental value etc. Although almost intensively cultured plant rightly comes under the domain of horticulture, primary effort is centered on the various traditional "garden" plants [6].

Lactuca sativa is an annual plant of the daisy family *Asteraceae*. It is most often grown as a leaf vegetable. Generally grown as hardy annual, lettuce is easily cultivated, although it requires relatively low temperatures to prevent it from flowering quickly. It can be plagued with numerous nutrient deficiencies, as well as insect and mammal pests and fungal and bacterial diseases [4-6]. *L. sativa* crosses easily within the species and with some other species within the *Lactuca* genus; although this trait can be a problem to home gardeners who attempt to save seeds, biologists have used it to broaden the gene pool of cultivated lettuce varieties. *L. sativa* is most often used in salads, although it is also used in the preparation of other kinds of food, such as soups, sandwiches and wraps; it can also be grilled. Assessment of heavy metal compositions in vegetable is one of the most important method used for monitoring environmental pollution [6], as the deficiency or excess of the metals is known to cause a number of serious metabolic disorders, as well as toxic effect [7]. The present study aimed at investigating the levels of some heavy metals (Pb, Zn, Ca, Cr and Cu), phytochemicals (Alkaloids, Flavonoids, Saponins, Tannins, Cardiac glycosides and Saponin Glycosides),

and proximate compositions (ash and moisture contents) of *L. sativa* grown in the vicinity of Cement Company of Northern Nigeria (Sokoto Cement) Sokoto.

2. MATERIALS AND METHODS

2.1 Chemicals and Reagents

The reagents used for the study included hydrochloric acid, trioxonitrate acid, sodium hydroxide, ferric chloride, Wagners reagent, sulphuric acid, Mayer's reagent, Fehling's solution, sodium chloride, perchloric acid, ammonium sulphate, chloroform, methanol, and distilled water. All other chemicals used were of analytical grade and purchased from standard manufactures.

2.2 Samples Collection

Fresh samples were collected in a pre-cleaned plastic bag from Kalambaina area Sokoto down

Cement Company of Northern Nigeria (Fig. 1) and Kwalkwalawa area around Usmanu Danfodiyo University Sokoto. The samples were identified and authenticated at the Department of Biological science, Usmanu Danfodiyo University, Sokoto. The collected samples were washed separately under running tap water before cut down into smaller pieces using knife and dried at 25-27°C for approximately 2 weeks. The samples were grounded into a fine powder, sieved through 2 mm sieve and stored in plastic jar labeled for analysis.

2.3 Extraction of Plant Samples

The procedure used was extraction by evaporation involving 5 g of the dried samples were put in a 250 ml conical flask and 100 ml of distilled water was added and was covered with Aluminium foil. It was allowed to stand for 72 hours and filtered using Whatman No.1 filter paper in a separate 250 ml conical flask. The filtrate was used for qualitative phytochemical analysis [8].



Fig. 1. Map of cement company of Northern Nigeria

2.4 Digestion Procedure

A 2.0 g of the samples were weighed into Kjeldahls flask mixed with 20 ml of concentrated sulphuric acid and helder tablet. The flask was heated at 70°C for about 40 min and then, the heat was increase to 120°C. The mixture turned to black after some time [9]. The digestion was completed after the solution became clear and white fumes appeared. The digest was diluted with 20 ml of distilled water and boiled for 15 min. Solution was then allowed to cool, it was transferred into 100 ml volumetric flasks and diluted to the mark with distilled water. The sample solution was then filtered through a Whatman filter paper No.1 (150 mm) into a screw capped polyethylene bottle, the procedure was repeated for all the samples.

2.5 Determination of Heavy Metals

The method applied for the assessments of heavy metal concentration of each sample of plant materials, after the digestion of the samples was by using the Atomic Absorption Spectrophotometric (AAS) techniques (AA320N) as described previously [10]. Each measurement was repeated 3 times, the mean and standard deviation were calculated (n=3).

2.6 Phytochemicals Screening

2.6.1 Alkaloids

The presence of alkaloids in each sample was investigated using the methods described by Wagner's [11]. 1 ml of each extract was treated with 2 drops of Wagner's reagent (2 g of iodine and 3 g of potassium iodine were dissolved in 20 ml of distilled water and made up to 100 ml with distilled water). Formation of brown precipitate indicates the presence of alkaloids in the extracts.

2.6.2 Flavanoids

The determination of the presence of flavanoids in the samples was done using alkaline reagent test by Okerulu et al. [12]. 3 ml of each extract were treated with 1 ml of 10% NaOH solution. Formation of intense yellow color, which becomes colorless on addition of dilute acid, indicates the presence of flavonoids in the extracts.

2.6.3 Saponins

The presence of saponins in the test samples was done using Harbone method [13] as

reported by Mercy [14]. 0.5 g of each extract was treated with 5 ml of distilled water and mixture was shaken vigorously, the production of foam which persisted in few minutes indicated the present of saponins in the extracts.

2.6.4 Tannins

The determination of the presence of tannins in the test sample was carried out using Ferric chloride test described by Harbone [12] as reported by Osagie [8]. 2 drops of 5% FeCl₃ was added to 1 ml of each extract. A greenish precipitate indicated the presence of tannins in the extracts.

2.6.5 Test for cardiac glycosides (Keller-Killiani Test)

1 ml of the filtrate was added to a test tube and then 2 ml of 3.5% FeCl₃ was added. The mixture was shaken for 1 minute and then 1 ml of concentrated H₂SO₄ was poured down the wall of the test tube so as to form a lower layer. A reddish brown ring at the interface indicates the presence of Cardiac Glycosides [12].

2.6.6 Test for Saponin glycosides

2 ml of the filtrate was added to a test tube, and then 2 ml of Fehling's solution was added. A bluish-green precipitate shows the presence of Saponin Glycosides [9].

2.6.7 Preparation of chloroform layer for Steroids and Anthraquinones

5 ml of the filtrate was added to a test tube, then 5 ml of chloroform was added, and the mixture was shaken vigorously for 1 minute. The mixture was allowed to settle until two layers are formed. The upper layer was separated from the lower layer (Chloroform layer) and discarded. The lower layer was used to test for Steroids and Anthraquinones [14].

2.6.7.1 Test for steroids (Lieber Mann Burchard Reaction)

1 ml of concentrated H₂SO₄ was added to 1 ml of the chloroform layer in a test tube and the mixture was allowed to settle. A reddish-brown ring at the interface indicates the presence of Steroids [6].

2.6.7.2 Test for Anthraquinones (Borntragers test)

5 ml of 10% NH₃ was added to the remaining chloroform layer in and shaken. The mixture was

allowed to settle and observation was made. A bright pink colour at the upper part of 2 layers formed indicates the presence of free Anthraquinones [12].

2.6.8 Test for volatile oils

2 ml of the filtrate was added to a test tube and 2 ml of 10% HCl was added. A white precipitation indicates the presence of Volatile oils [13].

2.6.9 Test for glycosides

1 ml of the filtrate was added to a test tube and 0.5mls of 50% H₂SO₄ for 15 minutes. After boiling, the mixture was allowed to cool and then it was neutralized with 1 ml of 10% NaOH. 2 ml of Fehlings' solution was added to the mixture and observation was made. A brick-red precipitate indicates the presence of glycosides [9].

2.7 Quantitative Phytochemical Analysis

2.7.1 Test for tannins

Powdered sample (0.1g) was put in a 100 ml conical flask and 52 ml volumetric flask. The residue was washed several times and the combined solution made up with distilled water to 0, 1, 2, 3, and 4 ml of the standard tannic acid and 10 ml of the sample solution in a 50 ml volumetric flask, 2.5 ml Folin-Denis reagent and 10 ml of Na₂CO₃ solution were added and made to volume with distilled water. The flask was allowed to stand for 20 minutes after which optical density was measured at 760 nm. The calibration curve was plotted from which the concentration of tannic acid in the sample was extrapolated [8].

2.7.2 Determination of alkaloid

5 g of powdered plant sample was extracted with 100 ml of methanoic: water (1:1; V: V) mixture and solvent evaporated. The resultant residue was mixed with 20 ml of 0.0025M H₂SO₄ and partitioned with ether to remove unwanted materials. The aqueous fraction was basified with strong NH₃ solution and then extracted with excess chloroform to obtain the alkaloids fraction or separated by filtration. The chloroform extraction was repeated several times and the extract was concentrated to dryness. The alkaloid was weighed and the percentage was calculated with reference to initial weight of powder [11].

$$\% \text{Alkaloid} = \frac{\text{Weight of alkaloid residue}}{\text{weight of sample}} \times 100$$

2.7.3 Determination of flavonoids

5 g of powdered sample was hydrolysed by boiling in 100 ml of hydrochloric acid solution for about 35 minutes. The hydrolysate was filtered to recover the extract (filtrate). The filtrate was treated with ethylacetate drop wise until in excess. The precipitated flavonoids were recovered by filtration using a weighed filter paper after drying in oven at 100°C for 30 minutes; it was cooled in a desiccator and reweighed. The difference in weighed gave the weighed of flavonoids which was expressed as the percentage of the weight of sample analysed.

$$\% \text{Flavonoid} = \frac{W_2 - W_1}{5g} \times 100$$

Where 5 g = weight of sample

W₁ = weight of empty filter paper

W₂ = weight of filter paper + sample precipitate

2.7.4 Determination of saponins

5 g of powdered sample was placed in a 250 ml conical flask containing 20 ml of 50% alcohol. The mixture was boiled under reflux for 30 minutes. The filtrate was allowed to cool at room temperature thereby resulting in the precipitation of saponins. The separated saponins were collected by decantation and suspended in about 2 ml of alcohol and filtered. The filter paper was immediately transferred to a desiccator containing anhydrous calcium chloride and the saponins were left to dry. They were weighed with reference of extract used.

$$\% \text{Saponins} = \frac{W_2 - W_1}{5g} \times 100$$

Where 5 g = weight of sample

W₁ = weight of filter paper

W₂ = weight of filter paper + sample precipitate.

2.8 Moisture Content Determination

Clean and dried petri-dishes were placed in an oven for about 30 minutes, cooled in a desiccator and each was weighed empty. 2 kg each of the dried samples were placed in a separate petri-dish and weighed. The petri-dishes and the content were then placed in an oven at 105°C for 24 hours. The percentage moisture of a given

sample was calculated using the formula in equation:

$$\%Moisture = \frac{W_2 - W_3}{W_2 - W_1} \times 100$$

Where W_1 = weight of empty petri-dish

W_2 = weight of petri-dish plus dried sample

W_3 = weight of petri-dish plus sample after drying [15].

2.9 Ash Content Determination

The term ash refers to the residue left after combustion of the oven dried sample and it is a measure of the total mineral content. The ash content of the samples was determined by heating porcelain crucibles in a muffle furnace to about 500°C, cooled in a desiccator and weighed. The 2 g each of the air-dried samples were transferred into separate crucibles and weighed. The crucibles containing the samples were placed in a muffle furnace and the temperature raised to 600°C for 3 hours. The crucibles containing the ashes were allowed to cool and weighed [10].

The percentage ash of a given sample was calculated using the formula in equation

$$\%Ash = \frac{W_3 - W_1}{W_2 - W_1} \times 100$$

Where W_1 = weight of empty crucible

W_2 = weight of crucible plus dried sample

W_3 = weight of crucible plus sample after drying

2.10 Determination of the pH of the Soil Samples

The process was carried out by weighing 20 g of each air-dried soil sample into separate 50 ml beakers. This was followed by addition of 20 ml of distilled water, which was allowed to stand for 30 minutes, stirring occasionally with a glass rod. The electrode of a pH meter already calibrated with pH 7.0 and pH 4.0 buffer solutions was dipped into each beaker containing the partly settled suspensions and readings were taken. The electrode was rinsed with distilled water and wiped dry with a clean filter paper each after reading [15].

2.11 Statistical Analysis

The collected data were subjected to statistical tests of significance using one way Anova

followed by dunnett comparison test ($P < 0.05$) to assess results in the lettuce samples. Probabilities less than 0.05 ($P < 0.05$) were considered significant. Statistical analyses were done using IBM SPSS (v20) statistical software. Results are Presented in Mean \pm Standard Deviation ($n=3$).

3. RESULTS AND DISCUSSION

Absorption of metals by plants in industrial areas may lead to phytotoxicity, the present study revealed significant difference ($P < 0.05$) in the metal concentration in plants found in industrial and non-industrial areas. The result showed a significant difference ($P < 0.05$) in moisture and ash contents of sample A (*Lactuca sativa* obtained from Kwalkwalawa) and sample B (*Lactuca sativa* obtained from Sokoto cement) (Table 1). The moisture and ash contents of sample A are low when compared with sample B. The values obtained are within the range of some vegetables grown in Nigeria [16-19].

Table 1. Moisture and Ash contents (%) in dry weight of *Lactuca sativa* samples

Sample	Moisture (%)	Ash (%)
A	45.33 \pm 1.33 ^a	35.5 \pm 2.18 ^a
B	13.30 \pm 1.31 ^b	19.9 \pm 1.02 ^b

Sample A = *Lactuca sativa* obtained from Kwalkwalawa
Sample B = *Lactuca sativa* obtained from Kalamaina industrial estate (Sokoto cement)

Superscript a&b in sample A and B respectively show significant difference ($P < 0.05$)

Phytochemicals are relevant in medicine, food, and dye industry. Some of them have pharmacological effects; for example, flavones, and tannins form important ingredients of several laxatives and medicine and in dyes. Qualitative estimation of phytochemicals of sample A and B revealed the presence of flavonoids, saponins, alkaloids, tannins, steroids, cardiac glycosides, and saponins glycosides (Table 2). This agrees with report by Miroslav and Vladimir [20]. Similarly, quantitative phytochemical analysis revealed presence of flavonoids, saponins, alkaloids, tannins, steroids, cardiac glycosides and saponin glycosides, in addition significant difference ($P < 0.05$) was observed between the two samples (Table 3).

Comparative studies of sample A and B showed significant difference ($P < 0.05$) in heavy metal (Pb, Cd, Zn and Fe) concentrations except Cr (Table 4). Pb has no biological importance; it is a

poisonous metal affecting almost every organ and system in the human body [21] and interferes with some metals such as Cd, Zn, and Fe. Prolonged exposure of Pb can lead to kidney damage. High Pb concentration in sample B (0.012 ± 0.002 mg/g) was observed when compared to its maximum acceptable concentration (MAC) in vegetables by WHO (0.002 mg/day) whereas no detection in sample A. Similar results were reported by Amusan *et al.* [22], Miller-Ihli and Baker [23] and contrary result was reported by Anthony and Balwant [24]. Cu is an essential element in trace amount and it plays a major role as co-enzyme in enzyme function, but in large amount it is toxic. Cu concentrations in sample A (0.005 ± 0.001 mg/g) and sample B (0.008 ± 0.002 mg/g) is within the acceptable limit of WHO (0.03 mg/day) for Cu. Cr in trivalent state (Cr^{3+}) is an essential trace element that potentiates insulin action and influences carbohydrate, lipid and protein metabolism. However, in hexavalent state (Cr^{6+}), it has toxic, mutagenic and carcinogenic effects [25]. The concentration of Cr in sample A (0.001 ± 0.001 mg/g) and B (0.002 ± 0.001 mg/g) are within the acceptable limit of WHO intake for Cr (0.002 - 0.005 mg/day).

Zn participates in the synthesis and degradation of carbohydrates, lipids, protein, and nucleic acids and has shown to play an essential role in polynucleotide transcription and translation and thus in the process of genetic expression. The concentrations of Zn of both samples are within the recommended values of WHO. Ca in sample A was found to be low (278.58 ± 8.230 mg/g) when compared to sample B (706.860 ± 14.980 mg/g). The latter is above the recommended limit by WHO 400 - 500 mg/g. The abnormal increase in Ca level might be as a result of cement deposition in Kalambaina area. Some of the studied results showed high values compared to other reports [2,26].

Table 2. Qualitative phytochemical analysis of *Lactuca sativa* samples

Phytochemical	A	B
Flavonoids	+	+
Saponins	+	+
Alkaloids	+	+
Anthraquinones	ND	ND
Tannins	+	+
Steroids	+	+
Volatile oils	ND	ND
Cardiac Glycosides	+	+
Saponin Glycosides	+	+

+ = Present

ND = Not detected

A = *Lactuca sativa* obtained from Kwalkwalwa

B = *Lactuca sativa* obtained from Kalambaina industrial estate (Sokoto cement)

The mineral analysis of soil collected from both sampled areas showed a significant difference ($P < 0.05$) in all metals with the highest concentrations observed in soil collected from Kalambaina (sample B). Low Pb concentration (0.002 ± 0.001 mg/g) was seen in sample A (soil sample from Kulkwalawa) when compared to (0.031 ± 0.002 mg/g) obtained from sample B (soil sample from Kalambaina) (Table 5). Similar results were reported with respect to industrialized areas [22-29]. The Cr concentration was observed to be 0.009 and 0.038 mg/g from soil analysis in sample A and B respectively which revealed high levels in sample B as well. Low value of Cr was reported by Uwah *et al.* [30]. The comparison of both soils showed significant difference in Zn and Cu as reported by Kachenko and Singh [31] but the value is low when compared with the values obtained during autumn in Torun Poland by Buszewski *et al.* [29] and also, the value of 0.102 mg/g by Amusan *et al.* [22,30,32].

Table 3. Quantitative phytochemicals estimation of the *Lactuca sativa* samples

Sample	Flavonoids	Saponins	Alkaloids	Tannins	Steroids	Saponin glycosides	Cardiac glycoside
A (%)	6.91 ± 0.15^a	5.04 ± 0.71^a	3.47 ± 0.96^a	2.29 ± 0.15^a	1.92 ± 0.45^a	4.10 ± 0.32^a	2.09 ± 0.62^a
B (%)	4.76 ± 0.32^b	2.65 ± 0.51^b	5.00 ± 0.27^b	0.30 ± 0.01^b	0.45 ± 0.36^b	1.71 ± 0.16^b	2.09 ± 0.58^b

Sample A = *Lactuca sativa* obtained from Kwalkwalwa

Sample B = *Lactuca sativa* obtained from Kalambaina industrial estate (Sokoto cement)

Superscript a&b in sample A and B respectively show significant difference ($P < 0.05$)

Table 4. Heavy Metals of *Lactuca sativa* collected from Kwalkwalawa and Kalambaina

Metals	Sample A (mg/g)	Sample B (mg/g)	WHO/FAO STD (mg/day)
Pb	ND	0.012±0.002 ^{bc}	0.002
Zn	0.008±0.002 ^a	0.043±0.003 ^{bc}	2.2
Ca	278.58±8.230 ^a	706.860±14.980 ^{bc}	400 – 500
Cr	0.001±0.001	0.002±0.001	0.002 - 0.005
Cu	0.005±0.001 ^a	0.008±0.002 ^b	0.03

Sample A = *Lactuca sativa* obtained from Kwalkwalawa

Sample B = *Lactuca sativa* obtained from Kalambaina industrial estate (Sokoto cement)

WHO/FAO values in mg/day are based on a 60kg body weight of adult

a = Comparison between A and WHO/FAO

b = Comparison between B and WHO/FAO

c = Comparison between Sample A and B

ND = Not Detected

Superscripts a, b&c in sample A, B and WHO/FAO STD respectively show significant difference (P<0.05)

Table 5. Heavy Metals of Soil(s) collected from Kwalkwalawa and Kalambaina

Sample	Pb (mg/g)	Zn (mg/g)	Ca (mg/g)	Cr (mg/g)	Cu (mg/g)
A	0.002±0.001 ^a	0.029±0.045 ^a	353.86±27 ^a	0.008±0.003 ^a	0.010±0.001 ^a
B	0.031±0.002 ^b	0.069±0.083 ^b	974.25±48 ^b	0.038±0.002 ^b	0.027±0.003 ^b

A = Soil obtained from Kwalkwalawa

B = Soil obtained from Kalambaina industrial estate (Sokoto cement)

Superscript a&b in A and B show statistically significant (P<0.05)

4. CONCLUSION

Our results revealed that the alkaline dust emitted from cement industry might be related to uptake of metals by the plants leading to decreased phytochemical concentrations. The level of Ca and Pb were found to be high in the study samples obtained from Kalambaina which exceeded the acceptable limit by WHO/FAO. This showed that the dust emitted by cement industries may play a role in accumulation of heavy metals in the environment, which may increase the possibilities of environmental pollution. However, dust depositions from cement industries; may not be the only cause of heavy metals accumulation, other activities such as; the excessive use of fertilizers, manures, pesticides, herbicides, and other agro-chemicals as well as the use of waste water by irrigating the soil may cause increase in the metal contents. Therefore vegetables grown within the vicinity of Northern Cement Company of Nigeria (Sokoto Cement) as food may not be safe for consumption at the time of study.

5. RECOMMENDATIONS

Further studies should be carried out to determine the concentration of heavy metals in water of the study areas. The study should also be extended to different areas of Kalambaina,

moverover, studies should also be carried out to assess heavy metals in blood of people residing at Northern Cement Company of Nigeria (Sokoto Cement) Kalambaina area.

ACKNOWLEDGEMENT

The authors are grateful to the Cement Company of Northern Nigeria (Sokoto Cement) for access to samples used for the study and Usmanu Danfodiyo University Sokoto where the research was conducted.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh SN, Rao DN. Growth of wheat plant exposed to cement dust pollution. Environ Poll (Ser. A). 1980;24:75-81.
2. Maina HM, Egila JN, Peter RA. Impact of cement dust deposition on plants in the vicinity of Ashaka cement factory, Gombe State, Nigeria. ARPN Journal of Science and Technology.2015;2225-7217.
3. Anthony EA, Esther UI. Levels of selected heavy metals in some Nigerian vegetables. Trends in Applied Sciences Research. 2007;2(1):76-79.

4. Kabata- Pendias A, Pendia H. Trace Elements in Soil and plants (3rd edition) CRS Press, Boca Raton Fla. London: CRC Press. 2000;413.
5. Abdulmojeed OL, Abdulrahman AA. Analysis of heavy metals found in vegetables from some cultivated irrigated gardens in the Kano metropolis, Nigeria. Journal of Environmental Chemistry and Ecotoxicology. 2011;3(6):142-148.
6. Bashiru I, Dalhat MH, Aminu AM, Sumayya AA, Zainab HB, Zulkallaini S. Determination of heavy metals and phytochemical analysis of some selected vegetables grown at Kalambaina Area Sokoto. Asian Journal of Soil Science and Plant Nutrition. 2017;2(2):1-7.
7. Agrawal. Effect of heavy metals contaminated vegetables. Indian Journal of Pharmacology. 2011;5:4-6.
8. Sofowora EA. Medicinal plants and traditional medicine in Africa. Spectrum Books Ltd, Ibadan, Nigeria. 2006;134-155.
9. Erwin JM, Ivo N. Determination of Lead in tissues: A pitfall due to wet digestion procedures in the presence of sulphuric acid. 1992;1(7):23-26.
10. A.O.A.C. Association of official analytical chemistry. 18th Edition, Washington DC. U.S.A; 1990.
11. Wagners. Method for detection of alkaloid in biological using wagners reagent (iodine in potassium iodide). International Epharmaceutica Scientia. 1991;1(1):357-377.
12. Okerulu IO, Onyema CT, Onwukeme VI, Ezech CM. Assessment of phytochemicals, proximate and elemental composition of *Pterocarpus soyauxii* (Oha) leaves. American Journal of Analytical Chemistry. 2017;80:406-415.
13. Harbone J. Experimental methods in analytical chemistry. New York, Chapman and Hall. 1973;140-145.
14. Mercy TA. Qualitative and Quantitative analysis of the phytochemicals in some selected species of the genus *Phyllanthus* in Zaria, Nigeria. 2015;710-711.
15. Daniel CH. Quantitative chemical analysis. Sixth Edition. W. H. Freeman Company, New York. 2003;364:132-145.
16. Hassan LG, Umar KJ, Usman A. Nutrient Content of leaves of *Tribulus terrestris* (Tsaida). Journal of Tropical Bioscience. 2005;5(2):77-82.
17. Asibey-Berko E, Tayie FAK. Proximate analysis of some Under Utilized Ghanaian vegetables. Ghana Journal of Science. 1999;39:91-92.
18. Nwaogu LA, Ujowundu CO, Mgbemena AI. Studies on the nutritional and phytochemical composition of *Amaranthus hybridus* leave. Journal of Bio Research. 2000;4(1):28-31.
19. Hassan LG, Umar KJ. Nutritional value of Balsam Apple (*Momordica balsamina* L.) Leaves. Journal of Nutrient. 2006;5(6): 522-529.
20. Miroslav R, Vladimir NB. Practical environmental analysis. Royal Society Chemistry. U.K. 1999;267-273.
21. FDA (Food and Drug Administration); U.S Supporting Document for Recommended Maximum Level for Lead in Candy Likely to be Consumed Frequently by Small Children.
22. Amusan AA, Ige VD, Oluwale R. Characteristics of soils and Crops uptake of Metals in Municipal Waste Dump Sites in Nigeria. J Hum. Ecol. 2005;17(3):167-171.
23. Miller-Ihli JN, Baker AS. Food and dairy products, applications of atomic spectroscopy. Encyclopaedia of Spectroscopy and Spectrometry. 2000;1: 583-588.
24. Anthony K, Balwant S. Heavy metals concentration of HOME grown vegetables near Metal Smelters in New South Wales Australia. 2006;2(1):76-79.
25. Guthrie AH. Introductory nutrition. Seventh edition, Times Mirrow/Mosby College Publishing. St. Louis. 1989;289-331.
26. Aksoy A, Sahin U, Duman F. Robina pseudo-acacia L as a possible biomonitor of heavy metal pollution. Turk J. Bot. 2000; 24:279-284.
27. Aiboni UV. Characterization and classification of soils of a representative topographic location in the University of Agriculture Abeokuta. ASSET Series A1. 2001;1(1):35-50.
28. Ibitoye AA, Ipinmoroti KO, Amoo IA. Effects of municipal refuse dump on the physico-chemical properties of Soil and Water. Nigerian Journal of Soil Science. 2005;15(2):122-128.
29. Buszewski B, Jastrzebska A, Kowalkowski T, Gorna-Binkul A. Monitoring of heavy metals uptake by plants and soils in the area of Torun, Poland. Polish Journal of Environmental Studies. 2000;9(6):1-7.
30. Uwah EI, Ndahia NP, Ogugbuaja VO. Study of the leaves of some agricultural

- pollutants in soils and water leaves (*Talinum triangulare*) obtained in Maiduguri, Nigeria. Journal of Applied Sciences in Environmental Sanitation. 2009;4(2):21-25.
31. Kachenko A, Singh B. Heavy metals concentration of home grown vegetables near metal smelters in New South Wales Australia. 2006;1-5.
32. Kashif SR, Akram M, Yaseen M, Ali S. Studies on heavy metals and their uptake by vegetables in Adjoining areas of Hudhara Drain in Lahore. Journal of soil and Environment. 2009;28(1):7-12.

© 2017 Dalhat et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/22485>