



Anti-Inflammatory and Antitussive Activity of Asthpadose, a Phytomedicine used in the Treatment of Asthma in Côte d'Ivoire

Baibo Guy Léonce ^{a*},
Ackah Jacques Auguste Alfred Bognan ^a,
Brahima Kouame Koffi Raoul ^a, Yaye Yapi Guillaume ^a
and Obouayeba Abba Pacôme ^a

^a Department of Biochemistry-Microbiology, UFR Agroforestry, Jean Lorougnon Guédé University, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. Authors BGL and AJAAB initiated this work. Author BGL carried out the manipulations, the results and wrote the manuscript. Author AJAAB supervised the analysis of the results and the writing of the article. Authors BKKR, YYG and OAP participated in writing of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajrb/2024/v14i4293>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117668>

Original Research Article

Received: 20/03/2024

Accepted: 23/05/2024

Published: 25/05/2024

ABSTRACT

Introduction: The aim of this study is to contribute to the valorisation of an anti-asthmatic phytomedicine (Asthpadose) traditionally used in Côte d'Ivoire.

Aims: Evaluate antiasthma's activity of the aqueous extract of Asthpadose (EAA) was.

*Corresponding author: E-mail: baguyleonce@gmail.com;

Cite as: Léonce, B. G., Bognan, A. J. A. A., Raoul, B. K. K., Guillaume, Y. Y., & Pacôme, O. A. (2024). Anti-Inflammatory and Antitussive Activity of Asthpadose, a Phytomedicine used in the Treatment of Asthma in Côte d'Ivoire. *Asian Journal of Research in Biochemistry*, 14(4), 49–59. <https://doi.org/10.9734/ajrb/2024/v14i4293>

Methodology: The antiasthma's activity of the aqueous extract of Asthpadose (EAA) was evaluated by the determining of its anti-inflammatory activity using the carrageenan oedema induction method. This anti-asthmatic activity was also evaluated by its antitussive action on the frequency of cough induced in rats exposed to liquid ammonia.

Results: The results revealed that EAA showed a very significant strong anti-inflammatory activity with an oedema inhibition rate ranging from 34.29% to 70.95%. This activity was much better than that of diclofenac sodium (a reference molecule against inflammation) with an oedema inhibition rate ranging from 12.24% to 58.39%. As for its antitussive activity, the results also showed good antitussive activity of the extract. EAA presented a cough inhibition rate of 74.22% compared to 58.76% for dextrometaphan which is also a reference molecule against cough.

Conclusion: This observed activity could therefore justify the use of Asthpadose in the treatment of asthma which is a disease manifested by inflammation and cough. In view of the very promising results of this study, it would therefore be necessary to continue studies on this phytochemistry to understand its mechanism of action on asthma.

Keywords: Asthpadose; rats; dextromethorphan; sodium diclofenac; ammonia; carrageenan.

ABBREVIATIONS

EAA : *Extrait Aqueux d' Asthpadose*

DFS : *Diclofenac sodium*

ED : *Distilled water*

DMT : *Dextrometorphane*

1. INTRODUCTION

Inflammation is a defense mechanism of the body against attacks of physical, chemical, biological or infectious origin [1,2]. This protective immune response can sometimes be harmful, linked to the pathogen and its persistence, to abnormalities in the regulation and production of pro-inflammatory cells [2,3]. As for coughing, it represents a means of sudden expulsion of air coming from the lungs thanks to the contraction of the expiratory muscles in order to rid the respiratory tract of foreign bodies and excessive secretions. Coughing also represents a defense reaction of the body and protection of the lung, which takes over when the muco-ciliary system which lines the respiratory tree no longer plays its role. It is a reflex act following irritation of the respiratory tract, the pleura or the diaphragm, which allows bronchial secretions (mucus) or a foreign body to be evacuated [4-6]. Inflammation and cough also appear during an asthma manifestation [7,8]. Asthma is a chronic inflammatory disease of the airways representing a global public health problem. This disease affects around 350 million people worldwide [9].

Asthpadose is a phytochemistry used in Côte d'Ivoire in the traditional treatment of asthma, allergy, liver disease and constipation.

The objective of this study is to evaluate the anti-asthmatic activity of Asthpadose. For the realization of this present project, a preparation of the aqueous extract was made from the Asthpadose recipe (Aqueous Asthpadose) then the anti-inflammatory and antitussive activities of Asthpadose were carried out on *Wistar* rats.

2. MATERIALS AND METHODS

2.1 Materials

2.1.1 Plant material

The plant material is a recipe called Asthpadose. This recipe is composed only of different parts (leaves, stems, flowers and roots) of medicinal plants. The plants that make up this recipe were harvested in Daloa. They were dried separately away from the sun for 2 weeks then sprayed individually. The different powders were weighed in proportional quantities then mixed to form Asthpadose.

2.1.2 Animal material

The animal material used in this study consists of male and female white rats of the *Wistar* albino strain. These animals come from the animal facility of the laboratory of the UFR of Pharmaceutical and Biological Sciences of the Félix Houphouët Boigny University. These animals were acclimated to room temperature and fed twice a day with IVOGRAIN granules then hydrated with tap water. These adult male and female albino rats were aged 3 to 4 months and weighed between 120 and 190 g and were then used to study the anti-inflammatory and antitussive activity.

2.2 Methods

2.2.1 Preparing the extract

The Asthpadose powder was used to prepare the aqueous extract of Asthpadose (EAA) which was obtained after pulverizing the different plants that constitute it. These plants were first collected in Daloa. They were washed, dried separately out of the sun for 2 weeks then sprayed individually. These different plants were then weighed in proportional quantities and then mixed. The preparation of the extract used during this study (aqueous extract of Asthpadose) was obtained according to the method of Zirihi *and al.*, [10,11]. For this preparation, 100 g grams of powder from the Asthpadose recipe was macerated in a liter of distilled water then homogenized in a blender. The homogenate obtained was successively filtered twice on hydrophilic cotton then on Whatman No. 3 filter paper. The filtrate obtained was dehydrated using an oven at a temperature of 55°C for 3 days. The dry evaporate was recovered on the 4th day in the form of a brown paste and constituted the aqueous extract of Asthpadose (EAA).

2.2.2 Preparation of animals

The animals were chosen at random and then marked to allow individual identification. Then, they were kept in their cages for acclimatization to laboratory conditions for 7 days before the experiment.

2.2.3 Anti-inflammatory activity of Asthpadose aqueous extract (EAA)

The anti-inflammatory activity test was carried out according to the method of Winter *et al.* [2,12,13] with some modifications. It was carried out by induction of oedema in the right hind leg of the rat using carrageenan (1%). 30 rats weighing between 120 g and 190 g were fasted for 12 h. These rats were divided into 5 groups of 6 rats each. The diameter of the leg at time T0 of each animal was determined using a caliper. The rats were distributed as follows: batch 1 (control batch) received distilled water, batch 2 received distilled water at a dose of 1ml/100g bw. Rats from batch 3 (reference control) received Sodium Diclofenac (25 mg/kg bw). As for batches 4 and 5 (test batches), they received EAA at respective doses of 250 mg and 500 mg/kg bw 1 hour after administration of distilled water, Diclofenac and EAA by gavage,

the 1% carrageenan, was injected cutaneously into the plantar aponeurosis of the right hind paw of each rat at a volume of 0.1 ml. The evolution of the oedema of the right paw was then determined at 1 h, 2 h, 3 h, 4 h, and 5 h [14] then the increase in oedema was assessed by determining the average percentage increase (%AUG) in the volume of the rat paw according to the formula:

$$\%AUG = ((V_t - V_0) / V_0) \times 100$$

% AUG: Percentage increase of the paw

V_t: Volume of the paw at time t

V₀: Initial volume of the paw [2,14]

The anti-inflammatory activity was finally evaluated by calculating the percentage of inhibition (% INH) of oedema according to the formula:

$$\%INH = ((\%AUG_{TM} - \%AUG_{TT}) / \%AUG_{TM}) \times 100$$

% INH: Percentage of oedema inhibition

%AUG_{TM}: percentage increase in cookies

%AUG_{TT}: percentage increase in Treaties [2,14]

2.2.4 Effect of Asthpadose extract on cough

The experiment was carried out according to the method described by [13,15,16] with some modifications. To make it, a grid under which a bowl has been placed will be placed in a cage. The grid served as a platform for the movement of animals exposed to the ammonia contained in the bowl. 3 mL of liquid ammonia (25% NH₄OH) are taken using a syringe then poured into the bowl. 30 seconds later, the animal was exposed to liquid ammonia for 2 minutes on this platform in the hermetically closed cage. The rat was taken out of this cage and placed in a second cage (observation cage). Each cough was detected by visual observation of the animal. The number of times the animal coughed was recorded over 5 minutes. Observation time was measured using a stopwatch.

2.2.4.1 Animal behavior

The detection of cough in animals placed in observation cages after exposure to liquid ammonia was described by Morice *et al.*, [13,17]. Cough was detected by the following symptoms: mouth opening, characteristic sound and postural changes (stretching of the front legs and forward stretching of the neck) accompanied by inhalation and exhalation.

2.2.4.2 Inhibition of cough frequency

The animals exposed to ammonia were randomly divided into 5 groups of 6 animals (3 males and 3 females). The animals from batch 1 (control batch) were untreated. The second batch (batch 2) received 1 mL/100 g bw of distilled water. As for batch 3 (reference control), the rats in this batch received Dextromethorphan (20 mg/kg bw). The rats in batches 4 and 5 received the Asthpadose extract at respective doses of 250 mg and 500 mg/kg of body weight with a quantity of 1 mL/100 g of body weight. The animals were then placed back in the cage. experiment (cage containing liquid ammonia 25 % NH₄OH) 1 hour after administration of extract and distilled water to be exposed again for 2 minutes. The number of coughs was again determined during the first 5 minutes.

The percentage of cough frequency was calculated by the following formula:

$$\% FT = (1 - T / C) \times 100$$

%FT: percentage of cough frequency

T: number of coughs recorded after treatment of animals with Asthpadose

DC: number of coughs emitted by the animals in the control batch (Lot 1) [13,18]

The percentage of cough inhibition was calculated by the following formula:

$$\% IT = (1 - Ta / Ca) \times 100$$

%IT: percentage of cough inhibition

Your: percentage of cough frequency of animals from batches treated with Asthpadose

That: Average diameter of the legs of animals from the control batch (Lot 1) [13,18]

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Action of asthpadose on inflammation

3.1.1.1 Measurement of oedema induced by carrageenan

The EAA results on the increase in paw oedema in rats treated with carrageenan are recorded in Table 1. These results showed an increase in paw diameter (oedema) over time in control animals as well as those of treated animals. During this study, no side effects or mortality

were observed in rats given carrageenan by injection.

The diameter of the paw of batch 1 animals (control batch) and batch 2 (treated with 2 ml/100g of distilled water) gradually increased by 0.20 ± 0.0 cm for batch 1 and by 0.19 ± 0.0 cm for batch 2 from the first hour to reach a maximum diameter at the fifth hour with a diameter increase of 0.30 ± 0.03 cm for batch 1 and 0.25 ± 0.03 cm for batch 2 while the paw diameter of rats treated with sodium diclofenac reaches its maximum increase in the first hour 0.19 ± 0.03 cm then gradually decreases from the second hour to the fifth hour 0.14 ± 0.07 cm.

However, in rats from batch 4, the batch having received 250 mg/kg bw of EAA, the oedema reached its maximum in the first hour with an increase in diameter of more than 0.17 ± 0.02 cm then was reduced from the second hour with a diameter of 0.16 ± 0.01 cm to reach a constant diameter of edema which was 0.16 ± 0.07 cm from the fourth hour until the fifth hour.

As for the rats treated with the dose of 500 mg/kg bw of EAA (batch 5), the oedemas in these rats reached their maximum increase at the first hour 0.15 ± 0.0 cm then decreased gradually from the second hour to fifth hour 0.10 ± 0.06 cm.

3.1.1.2 Percentage increase in oedema

The percentages of increase in paw oedema in rats are represented by Fig. 1. Indeed, until the 5th hour, the smallest value of the rate of increase was obtained in rats from batch 5 (treated batch with the dose 500 mg/kg bw of EAA). This value was 2.96 %. The rate of increase of batch 5 is followed by that of batch 3 (batch treated with 25 mg/kg bw of sodium diclofenac) with a slightly higher rate of increase. This rate is also followed by that of batch 4 (batch treated with the dose 250 mg/kg bw of EAA) with respective values of 4.24% and 7.24%. As for batch 2, its rate of increase remains the highest with a rate of 10.19 compared to the control batch but lower than those treated with diclofenac and EAA. The percentages of increase in oedema in batches 1 and batch 2 were statistically identical to each other ($p > 0.05$). On the other hand, the percentage increases in batches 3, 4 and 5 were statistically identical but lower than those of the control (batch 1) and batch 2. This difference in frequency compared to the control is very significant ($p \leq 0.01$).

Table 1. Effect of asthpadose extract on carrageenan-induced paw oedema

Treatment	0 hours	1 hour	2 hours	3 hours	4 hours	5 hours
Batch 1 (Control batch)	0	0.20 ± 0.0	0.24 ± 0.01	0.25 ± 0.06	0.28 ± 0.03	0.30 ± 0.03
Batch 2 (ED 2ml/100g)	0	0.190 ± 0.0	0.237 ± 0.04	0.25 ± 0.05	0.25 ± 0.03	0.25 ± 0.03
Batch3 (DFS 25)	0	0.19 ± 0.030	0.13 ± 0.027	0.13 ± 0.084	0.15 ± 0.072	0.14 ± 0.070
Batch4 (EA 250)	0	0.17 ± 0.02	0.16 ± 0.01	0.17 ± 0.06	0.16 ± 0.07	0.16 ± 0.02
Batch5 (EA 500)	0	0.15 ± 0.0	0.12 ± 0.05	0.12 ± 0.07	0.11 ± 0.07	0.10 ± 0.06

Batch 1: untreated control batch; batch 2 batch treated with distilled water; batch 3: batch treated with sodium diclofenac at a dose of 25 mg/kg bw; Batch 4 batch treated with Asthpadose extract at a dose of 250 mg/kg bw; batch 5 batch treated with Asthpadose extract at a dose of 500 mg/kg bw; ED: distilled water; DFS: diclofenac sodium; EA: Asthpadose extract




Fig. 1. Percentage increase in paw oedema induced by carrageenan

batch 1: untreated control batch; batch 2: batch treated with distilled water; batch 3: batch treated with sodium diclofenac at a dose of 25 mg/kg bw; batch 4: batch treated with Asthpadose extract at adose of 250 mg/kg bw; batch 5: batch treated with Asthpadose extract at a dose of 500 mg/kg bw; ED: distilled water; DFS: diclofenac sodium; EAA: Asthpadose extract

3.1.1.3 Percentage inhibition of edema

The percentages of inhibition of paw oedema in rats 1 hour after injection of carrageenan are presented in Fig. 2. This result revealed that from the first hour to the fifth hour, EAA at a dose of 500 mg /kg bw gave the highest percentage of inhibition of carrageenan-induced oedema with values of 34.29 % up to 70.95 %. This activity is followed by that of sodium diclofenac with a percentage of inhibition ranging from 12.24% to 58.39% then by EAA at

a dose of 250 mg/kg bw which presented an inhibition rate of 24.32 % to 55.64 %. The oedema inhibition percentages of batches 1 and 2 were statistically identical to each other ($p > 0.05$). Furthermore, those of batches 3, 4 and 5 were statistically identical ($p > 0.05$) but higher than those of the control (batch 1) and batch 2. This difference in frequency compared to the control was very significant ($p \leq 0.01$). At a dose of 500 mg/kg, EAA (unpurified extract) seemed to be more effective than sodium diclofenac, which is a reference molecule.




Fig. 2. Percentage of inhibition of paw oedema induced by carrageenan

Batch 1: untreated control batch; batch2: batch treated with distilled water; batch 3: batch treated with sodium diclofenac at a dose of 25 mg/kg bw; batch 4: batch treated with Asthpadose extract at a dose of 250 mg/kg bw; batch 5: batch treated with Asthpadose extract at a dose of 500 mg/kg bw; ED: distilled water; DFS: diclofenac sodium; EA: Asthpadose extract

3.1.2 Result of the action of asthpadose on cough

3.1.2.1 Influence of Asthpadose aqueous extract on the frequency of coughs caused by ammonia

The in vivo study of the effect of EAA on cough gave the results illustrated in Fig. 3. The results present the influence of EAA on the frequency of coughs caused by ammonia in rats. Untreated batch 1 indicated a cough frequency of 19.40 ± 0.93 . As for batch 2 (batch treated with distilled water), the frequency of coughing was 17.20 ± 0.86 ; while batch 3 which received 20 mg/kg bw

of dextromethorphan revealed a cough frequency of 8.0 ± 0.71 . The cough frequencies provided by the animals in batches 4 and 5 having received the respective doses of 250 and 500 mg/kg bw, were respectively 10.0 ± 1.41 and 5.0 ± 0.71 . The cough frequencies of batches 1 and batch 2 were statistically identical to each other ($p > 0.05$). Furthermore, the cough frequencies of batches 3, 4 and 5 were statistically identical ($p > 0.05$) but lower than those of the control (batch 1) and batch 2. This difference in frequency compared to the control was very significant ($p \leq 0.01$). However, the cough frequency value of batch 5 remained the lowest.




Fig. 3. Frequency of ammonia-induced cough

ED: distilled water; EAA: aqueous extract of Asthpadose; DMT: dextrometrophan L1: batch 1; L2:batch 2; L3: batch 3 L4: batch 4 and L5: batch 5 Values were expressed as means \pm SEM ($n = 6$). The mention ns and ** represents statistical significance. The urea values of the animals from the milked batches (L3, L4 and L5) compared to that of batch 1 were statistically very significant ($p \leq 0.01$).




Fig. 4. Percentage of cough inhibition

ED: distilled water; EAA: aqueous extract of Asthpadose; DMT: dextrometrophan L1: batch 1; L2: batch 2; L3: batch 3 L4: batch 4; and L5: batch 5 Values were expressed as means \pm SEM ($n = 6$). The mention ns and ** represents statistical significance. The urea values of the animals from the milked batches (L3, L4 and L5) compared to that of batch 1 were statistically very significant ($p \leq 0.01$).

3.1.2.2 Cough inhibition percentage

The evaluation of the antitussive activity of EAA is represented by Fig. 4. The result obtained with the animals of batch 2 to which distilled water was administered was 11.34 %. While the cough inhibition percentage of batch 3, batch having received 20 mg/kg bw of dextromethorphan, was 58.76 %. As for that of batch 4 (lot which received a dose of 250 mg/kg bw of EAA), it was 48.45 %. Finally, batch 5 which received the 500 mg/kg bw dose of the same extract gave an inhibition percentage of 74.22 %. Comparison of cough inhibition percentages showed statistically identical inhibition percentages ($p > 0.05$) between batch 3, batch 4 and batch 5. However, comparing batch 3, batch 4 and batch 5 to the control batch, the difference in the percentage of inhibition was statistically very significant ($p \leq 0.01$). Generally speaking, the percentage of inhibition of EAA at the dose 500 mg/kg bw remained the highest followed by dextromethorphan which is a reference molecule at 20 mg./kg bw then EAA at the dose 250 mg/kg bw to finish with distilled water.

3.2 Discussion

Inflammation is a reaction phenomenon implemented by the organism whenever the integrity of its morphological and biological constants is threatened [19]. The injection of carrageenan into animals causes local inflammation caused by tissue damage which results from the action of the prostaglandins and histamine produced. These mediators increase the permeability of capillaries in the region. As a result, exudate escapes from the bloodstream into the interstitial space. This exudate is the cause of localized oedema, which in turn compresses the nerve endings and thus causes a sensation of pain [20,21]. Carrageenan-induced paw oedema occurs in two phases. A phase involving the release of pro-inflammatory compounds such as histamine, serotonin and kinins which takes place in the first hour ($T = 1$ h) and a late phase due to an increase in COX-2 and release of PGE2 manifesting after the first hour ($T > 1$ h) [22,23]. According to Ghedira [24], the reduction in oedema of the aqueous extract of *Asthpadose* could be explained by a probable blockade of the excitation of the nociceptive afferent nerve endings of the extract. The activity anti-inflammatories linked to the presence of flavonoids in *Asthpadosis* was shown by

Baumann *and al.*, in [25,26]. The aqueous extract of *asthpadose* acts on inflammation by reducing prostaglandin. This result could be justified by the study carried out by Morimoto *and al.*, [27] who showed that flavonoids acted on prostaglandins, which are involved in the final phase of acute inflammation and pain. Uguru [28] and Hajjaj [29] have shown the effect of flavonoids in inhibiting the synthesis of prostaglandins and the effect of tannins in inhibiting arachidonic acid from anti-inflammation.

The study of the antitussive activity of *Asthpadose* consisted of inducing a dry cough by exposing rats to the vapor of a liquid ammonia solution (NH₂OH, 25%) and treating them with different doses of EAA in comparison with dextromethorphan, a reference cough suppressant. Liquid ammonia is a cough inducer which causes a reduction in respiratory amplitudes, this is bronchoconstriction. [30]. Irritation of RARS (Rapidly Adapting Receptors or irritant receptors) [31] causes contractions of the intercostal muscles. This irritation would be due to the release of histamine and the formation of other mediators [31,32]. Histamine causes bronchoconstriction, vasodilation and increased capillary permeability [33]. During this study, a significant reduction in the number of coughs in animals was observed after treatment with dextromethorphan as well as aqueous extracts of *Asthpadose*. The aqueous extract of *Asthpadose* could therefore have an action on the central nervous system, particularly at the level of the brainstem since, according to studies carried out by Cantekin *and al.* [34] and Gavliakova *and al.* [35] dextromethorphan would have a central action on the central nervous system at the brainstem level. It stimulates mu and kappa opioid receptors by depolarization of the vagus nerve. This depolarization results in an increase in the cough threshold, leading to a reduction in cough frequency. The aqueous extract of *Asthpadose* therefore demonstrated potential antitussive power in vivo. Also, Smith [36] showed that dextromethorphan acts on N-methyl-D-aspartate (NMDA) receptors in the central nervous system. These results would be comparable to those obtained in the study conducted by Jain *and al.* [18] on *Caesalpinia bonducella* in mice. These authors showed that extracts of this plant have antitussive activity. These results are also in agreement with those of Agnero [13] by evaluating the antitussive

activity of *chrysophyllum welwitschi* Engl. in rats and comparing it to that of dextromethorphan. Also, just like asthma, cough is caused by a stimulus and this stimulus would be inflammatory (allergens), mechanical, chemical (inhalation of gas, smoke) or thermal [37]. It is therefore probable that the aqueous extract of *Asthpadose* reduces the number of coughs by inhibiting either the inflammatory process or the nerve impulse responsible for triggering the cough [38]. Also, according to some researchers, cough is linked to mixed airway inflammation involving interactions between eosinophils and other cells such as neutrophils [39-42].

4. CONCLUSION

This study is part of the valorisation of medicinal plants and especially the search for new molecules from plant extracts. During this work, the aqueous extract of *Asthpadose* (EAA) used showed therapeutic action on inflammation and cough. Indeed, it would act on inflammation by inhibiting prostaglandins, histamine and arachidonic acid from inflammation. Also, its antitussive activity would be linked by its action on the central nervous system at the level of the brain stem. The beneficial effect of the *Asthpadose* tested would probably be attributable to the bioactive compounds it contains. These beneficial effects could also result from its combined action on prostaglandins, histamine and arachidonic acid in inflammation. However, additional studies will be necessary to understand the cellular and molecular mechanisms linked to these anti-inflammatory and antitussive activities.

ETHICAL APPROVAL

The study was approved by the Institutional Ethics Committee.

ACKNOWLEDGEMENTS

We would like to thank the Agrovalorization laboratory of the Agroforestry UFR of Jean Lorougnon Guédé University for providing us with the working framework. We also thank the traditional practitioners of Daloa (Côte d'Ivoire) for being collaborative in donating their phytomedicine for the completion of this work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Cheriti A, Rahmani S, Belboukhari N. Evaluation de l'activité anti-inflammatoire d'extraits aqueux de feuilles *Limoniastrum* feei (Plumbaginaceae). Algerian Journal of Arid Environment "AJAE". 2016;6(1):80-86.
2. Kouadio KJ, Ouattara-Soro FS, Abizi G, Zougrou NE, Kouakou KR, Begbin KE, Kplé TKM, Kablan KJJ, Koffi S. Activité anti-inflammatoire et études phytochimiques de l'extrait aqueux des écorces *Distemonanthus benthamianus* Baill. (Caesalpiniaceae Leguminosae - Caesalpinoideae). European Scientific Journal, ESJ. 2021;17(7):74.
3. Weill B, Bateux F, Dhainaut J. Immunopathologie et réactions inflammatoires. Eds, De Boeck, Université (Paris). 2003;12-23.
4. Debesse B, Rochemaure J. Dictionnaire de l'appareil respiratoire, CILF. Paris (France). 2008;336.
5. Dautzenberg B. Guide pratique de pneumologie. MMI. Paris (France). 2002; 336.
6. Aubaret C. La phytothérapie traditionnelle orientale et occidentale: convergences et différences dans le traitement de la toux. Thèse de Doctorat en Sciences Pharmaceutiques, Université Toulouse III Paul Sabatier, Toulouse, France. 2015;87.
7. Viladomiu, M, Hontecillas R, Bassaganya-Riera J. Modulation of inflammation and immunity by dietary conjugated linoleic acid. European journal of pharmacology. 2016; 785:87-95.
8. Polverino M, Polverino F, Fasolino M, Andò F, Alfieri A, De Blasio F. Anatomy and neuropathophysiology of the cough reflex arc. Multidiscip Respir Med. 2012;7 (1):1-5.
9. Berthe W. Ecole d'asthme et d'allergie d'Abidjan: opinion des participants. Thèse de Doctorat en Pharmacie et d'Odonto-Stomatologie, UNIVERSITE DE BAMAKO, Mali. 2010;113.
10. Zirihi G, Kra AKM, Guede-Guina F. Évaluation de l'activité antifongique de *Microglossa pyrifolia* (Lamarck) O. Kantze (Asteracée) « PYMI » sur la croissance in vitro de *Candida albicans*. Revue de médecine et pharmacie Afrique; 2003.
11. Yayé YG, Kra AKM, Ackah JAAB, Djaman AJ. Evaluation de l'activité antifongique

- etessai de purification des principes actifs des extraits de Terminaliamantaly (h.perrier), une combretacée, sur la croissance in vitro de Candida albicans. Bulletin de la Société Royale des Sciences de Liège. 2011; 80:953-964.
12. Winter CA, Risley FA, Nuss OW. Carrageenin induced oedema in hand paw of the rat as assays anti-inflammatory drugs. Experimental Biology Medicine. 1962;111:544-547.
 13. Agnero SM. Evaluation des activités anti-inflammatoire, antitussive, analgésique et antipyrétique de chrysophyllumwelwitschiiEngl. chez le rat et la souris. Thèse de doctorat de l'Université Félix HOUPHOUËT-BOIGNY, Spécialité Pharmacologie des Substances Naturelle. 2019;189.
 14. Anupama AS, Kishor NR, Rahul DK, Kanchan SM. Evaluation of anti-inflammatory and analgesic activities of Tamarindusindica seeds. International Journal of Pharmaceutical Sciences. 2012; 4(3):213-217.
 15. Xu SY, Bian RL, Chen X. PharmacologicalExperimentMethodology. People'sMedicalPublishing House, Beijing (China). 1991;1167.
 16. Yeo D, N'Guessan JD, Sea T, Coulibaly YA, Djaman AJ, Tako NA, Yavo JC, Guede- Guina F. Evaluation de l'activité antiasthmatique et antitussive de Combretum molle, plante medicinale de la pharmacopée ivoirienne. Phytotherapie. 2008;6(6):348-351.
 17. Morice AH, Menon MS, Mulrennan SA, Everett CF, Wright C, Jackson J,Thompson R. Opiate therapy in chroniccough. Am JRespirCrit Care Med. 2007;175:312-315.
 18. Jain S, Barik R, Yadav N, Singh S. Evaluation of anti- tussiveactivity of leaves of Caesalpinia bonducella. In experimentally induced cough in mice. IJPSR. 2013;4(1):415-418
 19. Singla AK; Pathak K. Tropical anti-inflammatory effets of Euphobia prostrate on carrageenan-induced foot pat oedema in mice. Journal of Ethnopharmacology. 1990;29:291-2994.
 20. Devulder PY, Hatron E, Hachulla E. Physiologie de l'inflammation. Cedex, Paris (France). 2002;15.
 21. Rousselet MC, Vignaud JM, Hofman P, Chatelet FP. Inflammation et pathologieinflammatoire. Paris Maloine. 2005; 320-331.
 22. Niu X, Li Y, Li W, Hu U, Yao H, Li H; Mu Q. The anti-inflammatory effects of Caraganatangutica ethyl acetate extract. J Ethnopharmacol. 2014;152(1):99-105.
 23. Lee SA, Moon SM, Choi YH, Han SH, Park BR, Choi MS, Kim JS, Kim YH, Kim DK, Kim CS. Aqueous extract of Codium fragile suppressed inflammatory responses in lipopolysaccharide-stimulated RAW264.7 cells and carrageenan-induced rats. Biomed Pharmacother. 2017; 93:1055-1064.
 24. Ghedira K. Les flavonoids: structure, propriétés biologiques, rôleprophylactique etemplois en thérapeutiques. Phytothérapie. 2005;4:162-169.
 25. Baumann K. Occupation alexposure to hexachlorocyclohexane. III. Neurophysiological findings and neuromuscular function in chronically exposed workers. International archives of occupational and environmenta lhealth. 1981;48:165-172.
 26. Emeraux E. Propriétés biologiques des flavonoïdes: étude bibliographique etévaluation de l'activité antioxydante. Thèse de Doctorat en Pharmacie, Université deLorraine, France. 2019;66.
 27. Morimoto A, Nakamori T, Watanabe T, Ono T; Murakami N. Pattern differences in experimental fevers induced by endotoxin, endogenous pyrogen, and prostaglandins. American Journal of Physiology. 1988;254:633-640.
 28. Uguru MO, Oluto PN; IOR D. Evaluation of analgesic and anti-inflammatory activities and phytochemical screening of the leaves extract of Paulliniapinnata (Sapindaceae). Journal of Chemical and Pharmaceutical Research. 2011;3(4):351-56.
 29. Hajjaj G. Screening phytochimique, etude toxicologique et valorisationpharmalogique de *matricariachamomilla* L. et de *l'ormenismixta* L. (asteraceae), rabat; 2017.
 30. Sanjay R, Dan VNJ, Beverly L; Mahdi M. Inhaled budesonide in the treatment of early COVID-19 (STOIC): a phase 2, open-label, randomised controlled trial. Lancet Respiratory Medicine. 2021;9:763–72.
 31. Widdicombe JG. Nervous receptor in the trachea bronchial tree. Program. of Brain Research. 1989;67:49–64.

32. Yu J, Zhang JF, Robertts AM, Collins LC; Flether EC. Pulmonary rapidly adapting receptor stimulation does not increase airway resistance in anesrhetized rabbit. American Journal of Respiratory and Critical Care Medicine. 1999;160:906-912.
33. Sudo T, Fumiaki H, Takashi N. Responses of trachea bronchial receptors to inhaled furosemide in anesthetized rats. American Journal of Respiration and Critical Care Medicine. 2000;162: 971-975.
34. Cantekin EI, Mandell EM, Bluestone CD. Lack of efficacy of a decongestant-antihistamine combination for otitis media with effusion ("secretory" otitis media) in children: results of a double-blind randomized trial. N Engl J Med. 1983; 308(6):297-301.
35. Gavliakova S, Biringeroza Z, Buday T, Brozmanova M, Calkovsky V, Poljacek I, Plevkova J. Antitussive effects of nasal thymol challenges in healthy volunteers. Respir Physiol Neurobiol. 2013;187(1): 104-107.
36. Smith JA, Woodcock A. Chronic cough. The New England journal of medicine. 2016;375:1544-1551.
37. Charpin D. Définition et épidémiologie de l'asthme. Poumon - plèvre - médiastin, EMC, Paris (France). 1984; 6039.
38. Harborne AJ. Phytochemical methods a guide to modern techniques of plant analysis. Springer Dordrecht, 3eme édition, Dordrecht (Pays- Bas). 1998;302.
39. Niimi A, Torrego A, Nicholson AG, Cosio BG, Oates TB, Chung KF. Nature of airway inflammation and remodeling in chroniccough. The Journal of allergy and clinicalimmunology. 2005;116:565-570.
40. Cepuc IA. La modulation du réflexe de toux par l'exercice chez le lapin sensibilise a l'ovalbumine. Thèse de doctorat de l'Université de Lorraine, Ecole Doctorale BioSE (Biologie-Santé-Environnement). 2017;120.
41. Adesina SK, Johnny II, Olayiwola G. Plants in Respiratory Disorders II- Antitussives, A Review. J. Pharm. Res. Int. [Internet]. 2017 May 5 [cited 2024 May 15];16(3): 1-21. Available:https://journaljpri.com/index.php/JPRI/article/view/219
42. Gandhi GR, de Sousa Leao GC, da Silva Calisto VK, Vasconcelos AB, Almeida ML, Quintans JD, Barreto E, Narain N, Júnior LJ, Gurgel RQ. Modulation of interleukin expression by medicinal plants and their secondary metabolites: a systematic review on anti-asthmatic and immunopharmacological mechanisms. Phytomedicine. 2020;70:153229.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/117668>