



Adie's Pupil (About a Case)

**M. Bouchaar^{1*}, S. Bajjouk¹, S. Haddougui¹, B. Houda¹, F. El Asri¹, K. Reda¹
and A. Oubaz¹**

¹*Department of Ophthalmology of the Cheikh Khalifa, Ibn Zaid University Hospital, Casablanca, Faculty of Medicine, Mohammed VI University of Health Sciences (UM6SS), Casablanca, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. Author MB designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors SB and SH managed the analyses of the study. Author BH managed the literature searches. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Panagiotis Tsikripis, University of Athens, Greece.

Reviewers:

(1) Waleed K. Abdulsahib, Al Farahidi University, Iraq.

(2) Shreya Thatte, Sri Aurobindo Medical College and PG Institute, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/64331>

Case Study

Received 20 November 2020

Accepted 28 December 2020

Published 20 January 2021

ABSTRACT

Adie syndrome is a relatively common neurological disorder of unknown etiology comprising unilateral or bilateral tonically dilated pupils with light-near dissociation and tendon are flexia . Adie syndrome is mostly idiopathic with no identifiable cause but may rarely be caused by local disorders involving the orbit that affect the ciliary ganglion including infections such as syphilis, varicella, autoimmune hepatitis, sarcoidosis. Adie's syndrome is a clinical diagnosis. Low-concentration pilocarpine (one-eighth to one-tenth percent) test may be useful to demonstrate the cholinergic denervation supersensitivity (80% prevalent) in the tonic pupil. The usual treatment of a standardised Adie syndrome is to prescribe reading glasses to correct for impairment of the eye(s) if it exist. Adie's syndrome is not life-threatening or disabling. As such, there is no mortality rate relating to the condition. We report a case of a young man without a history of medical pathology, who presented with an isolated pupil of adie for ten months.

Keywords: *Adie syndrome; neurological disorder; tendon; pupil; vision; oculomotor; muscles; ciliary ganglion; lyme disease; sjögren's syndrome; choroidal tumors.*

*Corresponding author: E-mail: bouchaar.m@gmail.com;

1. INTRODUCTION

Adie's syndrome is a relatively common neurological disorder of unknown etiology comprising dilated unilateral or bilateral pupils with near-lumen dissociation [1]. It is named after the British neurologists William John Adie and Gordon Morgan Holmes, who independently described the same disease in 1931 [2]. The syndrome is caused by damage to the post ganglion fibers of the parasympathetic innervation of the eye. It is frequently observed in women with, in particular, tendon areflexia [3]. Adie's syndrome is a clinical diagnosis. Low-concentration pilocarpine (one-eighth to one-tenth percent) test may be useful to demonstrate the cholinergic denervation supersensitivity (80% prevalent) in the tonic pupil [3,4]. After the administration of the dilute pilocarpine, a more miotic response will be seen in the affected eye compared with the normal fellow eye since these low concentrations are usually ineffective in normal pupils [4]. The tonic pupil may become smaller (miotic) over time which is referred to as "little old Adie's" [5]. We are reporting the case of a young man with an isolated Adie pupil.

2. OBSERVATION

A young man aged 27, with no particular history, presented to the Ophthalmology Outpatient Department with anisocoria discovered 10 months ago by the patient himself; without other accompanying signs.

On examination, he had a vision of 10/10 in both eyes. The photomotor reflex examination found a right pupil in reactive semi-mydriasis, the consensual reflex is present, there is no PDAR (Fig. 1). There was also no limitation of the oculomotor muscles. Eye tone and anterior

segment examination were normal in both eyes. The fundus examination was unremarkable. The examination of the osteotendinous reflexes was normal and the cerebral orbital MRI angiography is without abnormality. The pilocarpine test diluted to 0.125 is positive (Fig. 2). The remainder of the neurological, cardiovascular and osteotendinous examination was normal. So we concluded that it is a pupil of Adie.

3. DISCUSSION

Adie syndrome is primarily idiopathic with no identifiable cause, but can rarely be caused by local disorders involving the orbit that affect the ciliary ganglion, including infections such as syphilis, chickenpox, human parvovirus B19, the virus human immunodeficiency and Lyme disease [6,7], ischemia due to lymphomatoid granulomatosis, migraine and giant cell arteritis, [7,8] autoimmune diseases such as Sjögren's syndrome, polyarteritis nodosa, sarcoidosis, amyloidosis, Guillain-Barré and Vogt-Koyanagi syndrome - Harada disease [8], local or general anesthesia [9], orbital or choroidal tumors [10], orbital surgery for orbital floor fractures [11], neuromuscular conditions such as Lambert-Eaton myasthenic syndrome and paraneoplastic in association with anti-Hu antibodies [12]. It can also be caused by retinal photocoagulation which results in damage to the ciliary nerves in the suprachoroidal space [13].

The incidence of Adie's syndrome is approximately 4.7 / 100,000 inhabitants / year with a prevalence of two cases / 1,000 inhabitants (approximately) [14]. Young adults generally aged 25 to 45 are the most frequently affected, as is the case with our patient. On the other hand, there is a female predominance unlike our case, it was a male patient.



Fig. 1. Right eye showing a dilated pupil despite bright lighting (adie's pupil)



Fig. 2. Miosis pupil after the pilocarpine test confirming the diagnosis of the adie's pupil

West and Barnett [15] had reported ocular findings such as lens opacities, vitreous frosting, arteriosclerotic changes, eyelid abnormalities (stiffness or tightness in, telangiectasia), deficient tear secretion, and conjunctival abnormalities (injection and vascular sludging) in thirty-eight patients with systemic sclerosis. Anand [16] has described a patient with eyelid tightness and colloid bodies in retina.

Adie's syndrome is a clinical diagnosis. A low concentration pilocarpine test (one-eighth to one-tenth percent) may be useful in demonstrating supersensitivity to cholinergic denervation (80% prevalent) in the tonic pupil [17]. After administration of diluted pilocarpine, a more miotic response will be observed in the affected eye compared to the other normal eye since these low concentrations are generally ineffective in normal pupils [18].

There is no treatment and the pilocarpine, which contracts the pupil, is generally poorly tolerated; however, wearing tinted glasses does reduce photophobia [19].

We often see a stability of mydriasis, sometimes it decreases slightly over time. Sometimes we can see bilateralization (rare) [20].

4. CONCLUSION

Adie's pupil is a rare manifestation, the eye examination is important and the pilocarpine test remains the fundamental diagnostic element.

CONSENT

As per international standard or university standard, patient's written consent has been collected and preserved by the authors.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Holmes-Adie syndrome information page. National Institute of Neurological Disorders and Stroke. Archived from the original on 2007-10-16. Retrieved. 2008-01-21.
2. Siddiqui AA, Clarke JC, Grzybowski A. (November). "William John Adie: The man behind the syndrome" (PDF). *Clinical & Experimental Ophthalmology*. 2014;42(8):778–84.
3. Leavitt JA, Wayman LL, Hodge DO, Brubaker RF. Pupillary response to four concentrations of pilocarpine in normal subjects: Application to testing for Adie tonic pupil. *Am J Ophthalmol*. 2002;133(3):333-6.
4. ROSS AT. Progressive selective sudomotor denervation; a case with coexisting adie's syndrome. *Neurology*. 1958;8(11):809-17.
5. Weller M, Wilhelm H, Sommer N, Dichgans J, Wiethölter H. Tonic pupil, areflexia and segmental anhidrosis: Two additional cases of ross syndrome and review of the literature. *J Neurol*. 1992;239(4):231-4.
6. Sakai T, Shikishima K, Mizobuchi T, Yoshida M, Kitahara K. Bilateral tonic pupils associated with neurosyphilis. *Jpn J Ophthalmol*. 2003;47(4):368-71.

7. Stricker RB, Winger EE. Holmes-Adie syndrome and lyme disease. *Lancet*. 2001;357(9258):805. [PubMed].
8. Cerny R, Rozsypal H, Kozner P, Machala L. Bilateral Holmes-Adie syndrome as an early manifestation of the HIV neuropathy. *Neurol Sci*. 2010;31(5):661-3.
9. Foroozan R, Buono LM, Savino PJ, Sergott RC. Tonic pupils from giant cell arteritis. *Br J Ophthalmol*. 2003;87(4):510-2.
10. Purvin VA. Adie's tonic pupil secondary to migraine. *J Neuroophthalmol*. 1995;15(1):43-4.
11. Bennett JL, Pelak VA, Mourelatos Z, Bird S, Galetta SL. Acute sensorimotor polyneuropathy with tonic pupils and an abduction deficit: An unusual presentation of polyarteritis nodosa. *Surv Ophthalmol*. 1999;43(4):341-4.
12. Garza Leon M, Herrera-Jimenez IP, González-Madrigal PM. Complete Vogt-Koyanagi-Harada disease and Holmes-Adie syndrome: Case Report. *Ocul Immunol Inflamm*. 2014;22(4):336-40.
13. Guaraldi P, Mathias CJ. Progression of cardiovascular autonomic dysfunction in Holmes-Adie syndrome. *J Neurol Neurosurg Psychiatry*. 2011;82(9):1046-9.
14. Kobayashi M, Takenami T, Kimotsuki H, Mukuno K, Hoka S. Adie syndrome associated with general anesthesia. *Can J Anaesth*. 2008;55(2):130-1.
15. Goldstein SM, Liu GT, Edmond JC, Katowitz JA, Rorke LB. Orbital neuroglial hamartoma associated with a congenital tonic pupil. *J AAPOS*. 2002;6(1):54-5.
16. Bruno MK, Winterkorn JM, Edgar MA, Kamal A, Stübgen JP. Unilateral Adie pupil as sole ophthalmic sign of anti-Hu paraneoplastic syndrome. *J Neuroophthalmol*. 2000;20(4):248-9.
17. Wirtz PW, de Keizer RJ, de Visser M, Wintzen AR, Verschuuren JJ. Tonic pupils in Lambert-Eaton myasthenic syndrome. *Muscle Nerve*. 2001;24(3):444-5.
18. Kaur S, Aggarwal P, Jindal N, Dayal S, Jairath V, Jain VK, Viridi S. Harlequin syndrome: A mask of rare dysautonomic syndromes. *Dermatol Online J*. 2015;21(4).
19. Cheshire WP, Goldstein DS. The physical examination as a window into autonomic disorders. *Clin Auton Res*. 2018;28(1):23-33.
20. Zhang L, Luo S, Jin H, Lv X, Chen J. Anti-hu antibody-associated adie's pupil and paraneoplastic sensorimotor polyneuropathy caused by primary mediastinal small cell carcinoma. *Front Neurol*. 2019;10:1236.

© 2021 Bouchaar et al; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/64331>