



Nutritional Quality Differential, Growth and Economics Efficiency of Some Selected Commercial Floating Fish Feeds in Saki West Oyo State Nigeria

M. A. Adedokun^{1*}, Z. O. Oluwafemi², A. O. Ayanboye³ and G. Oladipupo²

¹*Department of Animal Health and Production Technology, The Oke-Ogun Polytechnic Saki, Nigeria.*

²*Department of Agricultural Economics, Oyo State College of Agriculture and Technology, Igbo-Ora, Nigeria.*

³*Department of Fisheries and Wildlife Management, Osun State University, Osogbo, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author MAA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors ZOO and AOA managed the analyses of the study. Author GO managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2019/v4i130043

Editor(s):

(1) Dr. Eugene A. Silow, Chair of Invertebrate Zoology and Aquatic Ecology, Institute of Biology, Irkutsk State University, Russia.

(2) Dr. Matheus Ramalho de Lima, Professor, Federal University of South of Bahia, Brazil.

Reviewers:

(1) Rakpong Petkam, Khon Kaen University, Thailand.

(2) Felix Eze, Nigeria Maritime University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50215>

Original Research Article

Received 15 May 2019
Accepted 26 July 2019
Published 10 August 2019

ABSTRACT

Nothing is more important than quality nutrition and adequate feeding of fish in captivity. Undernourished fish in terms of nutrient deficiency, cannot maintain its health for proportionate growth regardless the intense of feeding and quality of the environment.

A 49 day-feeding trial was carried out to investigate nutritional quality differential, growth and economics efficiency of some sampled commercially extruded floating feeds, based on frequency of usage among fish farmers in the study area. The sample feeds were sourced from respective distributors covering the zone of study. The feeds were designated as Fd₁, Fd₂, Fd₃, Fd₄, Fd₅ and Fd₆(control), with 3 replicates for each treatment. The examined growth performance, feed utilization and economic efficiency of feeds followed particular trend pattern and significantly

*Corresponding author: Email: matdokun@gmail.com, adediseas@gmail.com;

different ($p < 0.05$) across the sampled feeds (FW, MWG, SGR, TPI, PER and FCR). Finally, the control diet was least consumed, sustained positive allometry growth pattern and concomitant marginal profits; than feed 1 and 2 which had sharp drops in the growth pattern of fish after four (4) weeks.

Keywords: Allometry growth; economic efficiency; fish farming; feed utilization; differential; nutrition.

1. INTRODUCTION

Aquaculture is one of the fastest animal based food producing sectors, particularly in developing countries. However, success in aquaculture depends on the ability of a farmer to cost effectively meet the nutritional demand of the cultured fish species. This is because feed type as well as feed quality may have consequences on both growth efficiency and feed utilization [1]. Good nutrition in animal production system is essential to economically produce a healthy and high quality product. In fish farming, nutrition is critical because feed represents 50-60% of the production costs, [2]. The development of new species, specific diet support the aquaculture (fish farming) industry as it expands to satisfy increasing demand for affordable, safe and high quality fish.

As aquaculture production become more and more intensive in Nigeria, fish feed will be a significant factor in increasing the productivity and profitability of aquaculture [3]. The need to intensify the culture of the fish, so as to meet the ever increasing demand for fish has made it essential to develop suitable diet either in supplementary form for ponds or as complete feed in tanks [4]. The contribution of fisheries to the national economy is very significant in term of employment, income generation, poverty alleviation, food security, foreign exchange earnings and provision of raw materials for the animal feed industry [5]. Catfish (*Clarias* sp.) are the major commercially species in Nigeria, for good market and culture (management) reasons [6].

Since 2000 there has been a rapid expansion in urban aquaculture and a significant development in high density catfish culture. As a result of this intensification in catfish culture, the aqua feed industry has grown and concerted effort has been focusing on research in fish nutrition and fish diet which start at Nigeria Institute for Oceanography and Marine Research (NIOMR). Laboratory size pellet mill and about 12 commercial aqua feed producers were established in Nigeria to complement companies

that import high quality floating feed [7,8]. As such, there is currently in the market assortment of both imported and locally manufactured pelleted floating catfish feed brands.

Feed trial studies have been carried out on *Clarias gariepinus* to evaluate their growth response to different readily available protein sources [9,10,11,12,13,14,15]. The submission of their findings were not conclusive and the variations in conclusion of these afore-listed authors is a source motivation to further expand the body of knowledge regarding the nutritional quality differential, growth and economics efficiency of some selected floating feeds.

2. MATERIALS AND METHODS

2.1 Study Area / Experimental Site

35 practicing fish farmers in Saki – West L/Gvt. were sampled based on their preferred floating fish feeds abound in the market. The experiment was carried out in the Fisheries Laboratory of Oyo State College of Agriculture and Technology Igbo-ora, Oyo State, Nigeria.

2.2 Experimental Feeds (Sampled Feeds)

There are diverse of fish feeds which offer fish farmers' opportunity to choose out of the available floating fish feeds in the market. In this experiment, only five types of fish feeds were chosen based on frequency of usage by sampled fish farmers. They include, CF, AF, RF, VF, DF and CLF (Smoked fish waste meal) designated as Fd₁, Fd₂, Fd₃, Fd₄, Fd₅ and Fd₆ (control). The control diet was formulated using smoked fish waste (smoked fish waste meal) while other feed stuffs were purchased.

2.3 Experimental Design

The experimental design was in triplicates of eighteen plastic bowls for a period of 7 weeks, (April to June, 2018). Level of water in each bowl (60 cm x 30 cm x 15 cm) was maintained at 30 litres and the renewal of water was weekly (every seven days) at ratio 1:1 to avoid the shock as a

result of the seemingly new environment when the whole water is changed.

2.4 Experimental Procedure

One hundred and eighty (180) African catfish (*Clarias gariepinus*) juveniles of average weight 38.8 g, was obtained from a reliable source. The fish were acclimatized for 2 days and were fed with control diet feed (Fd_6) at 5% of their body weight twice daily; morning and evening. The fish were starved for 24 hours in order to empty their stomach and prepared their appetite for the new feed trials. The feeding ration and diet per meal were prepared at 5% body weight and two time feeding regimes. Weekly adjustment of feeding ration and diet were carried out throughout the experiment.

2.5 Growth Performance Parameters of Test Organism Juveniles Fed Different Feeds

Data on growth performance were collected weekly using the following nutrient utilization and growth parameters:

Main Weight gain = Final weight – initial weight

Average weight = Total weight / No of fish

Specific growth rate = $(\ln \text{ final body weight} - \ln \text{ initial body weight} / \text{Time (days)}) \times 100$

ADWG = MWG / Period of the experiment

%WG = MWG / Initial mean weight x 100

Protein Gain = MTPI g / Culture Time (days)

Total Protein Intake (TOi) = Total feed consumed x % CP in the feed

PER = Net weight gain (g) / Amount of protein fed (g)

Feed Conversion Ratio (FCR) = Total feed intake / Total wet weight gain

Total fish production: (Final weight g x Survival rate /1000)

2.6 Statistical Analysis of Data

One-way analysis of variance (ANOVA) was used to determine the effects of diets on growth

and nutrient utilization indices using 16,0 version of [16] statistical package. Significant differences between individual means were identified using the Duncan's multiple range test (Duncan, 1955). Mean differences were considered significant at $p < 0.05$.

2.7 Water Quality Management

The water quality variables such as Temperature, Hydrogen- Ion Concentration (pH) were measured with a combined digital pen-type daily meter, while dissolved oxygen (mg/l) was measured using Winkler's method and conductivity by a digit conductivity meter [17].

3. RESULTS AND DISCUSSION

Differential nutritional quality, growth Response and economic efficiency.

Nitrogen Free Extract (NFE) = $100 - (\text{Crude Protein} + \text{Crude lipid} + \text{crude fibre} + \text{total ash})$. Gross energy: Caloric value of protein 5.65, NFE 4.1 and lipid 9.45 kcal g^{-1} , Digestible energy: caloric value of protein 3.5, NFE 2.5 and lipid 8.1 kcal g^{-1} [18] (Adedokun et al., 2017).

3.1 Water Quality Parameters (WQP)

The mean water quality of the plastic trough system at weekly intervals during the study is presented in Table 2. Throughout the feeding trials, the water quality was keenly monitored. The observed water quality parameters were within the acceptable ranges of APHA/AWWA/WPCF [17,19].

The amounts of sampled feeds consumed were compared with weight gained by the fish. The graph shows wide variation in the sampled feed consumed (Fig. 1a) but relative little variation in the body weight gained (Fig. 1c) by the fish sampled. The wide range between quantity of feed consumed and weight gained ratio is shown in Fig. 1(b).

Acceptability and palatability of feed is a function of the processing methods.

The feed conversion ratio (FCR) depends on many factors such as feed palatability, fish breed and species, energy content, level of fibre inclusion, crude protein content, mineralization etc.

Table 1. Average mean values of proximate composition in experimental feeds

Parameters	DT1	DT2	DT3	DT4	DT5	DT6	Mean	SD
Crude Protein	28.70	50.75	27.65	35.35	29.05	40.02	35.25	8.97
Ash	6.17	4.02	5.26	5.50	5.90	6.28	5.52	0.83
Crude Fiber	0.01	0.01	0.02	0.01	0.01	0.35	0.7	0.35
Lipid Ether Extract	6.50	7.50	6.70	7.10	6.80	4.65	6.54	0.99
Moisture	91.99	92.67	92.39	91.70	92.15	93.76	92.44	0.73
Dry Mater								
NFE	58.62	37.72	60.37	52.04	58.24	48.7	52.62	8.55
Gross Energy (kcalg ⁻¹)	463.92	512.26	467.05	480.18	467.16	389.72	463.38	40.31
Digestible Energy (kcalg ⁻¹)	361.27	370.39	362.34	363.38	360.60	348.18	361.03	7.21
Energy/Protein ratio	12.6	7.3	13.1	10.3	12.4	8.7	10.73	2.37

Table 2. Water quality parameters of the experiment

Parameters	Dietary sampled feeds						Mean	SD
	Fd ₁	Fd ₂	Fd ₃	Fd ₄	Fd ₅	Fd ₆		
Temperature	27.58	27.57	27.46	27.40	27.40	27.46	27.48	0.08
DO (mg/l)	6.20	5.40	5.20	5.40	5.10	4.80	5.35	0.47
P ^H	6.40	6.80	6.70	6.60	6.60	6.60	6.62	0.13
Conductivity (µmoh/cm ³)	580	582	580	583	585	420	555	66.16

Table 3. Growth response efficiency of *Clarias gariepinus* juveniles

Parameters	DT1	DT2	DT3	DT4	DT5	DT6	Mean	SD
Total Feed	141.5	153.2	116.0	141.3	130.0	72.6	125.77	28.92
Final Weight	899.5	973.8	581.6	878.6	881	339.4	758.98	246.00
% Survival	80	86.6	66.6	86.6	86.6	90	82.80	8.55
% Mortality	20	13.3	33.3	13.3	13.3	10	17.20	8.54
Average Wt.	37.5	37.5	29.1	33.8	33.9	12.6	30.73	9.41
MWG	860.7	935	542.8	839.8	842.2	300.6	720.18	246.01
ADWG	17.6	19.1	11.1	17.1	17.2	6.1	14.70	5.03
% WG	2218.3	2409.9	1399	2164.4	2170.6	774.7	1856.13	634.05
SGR	17.6	19.1	11.1	17.1	17.2	6.1	14.70	5.03
TPI	40.6	77.8	32.1	50.0	37.8	29.1	44.57	17.83
PER	22.2	12.5	18.1	17.6	23.3	11.7	17.57	4.79
Total Fish Produced	21.6	25.3	11.6	22.8	22.9	9.2	18.90	6.73
Feed CR	0.16	0.16	0.20	0.16	0.14	0.22	0.17	0.30

Figs. 2a and 2b show the pattern of feed conversion ratio with different feeds sampled. The descriptive analysis revealed that diet 6 has highest feed conversion ratio (FCR) though less than 0.5 of total dry feed consumed divided by the wet weight of fish harvested. While diet 5 had the lowest feed conversion ratio. Theoretically, among other

factors that affect utilization nutrient include digestible protein content of the feed, energy-protein ratio, mineral and vitamin. Moreover, it was observed that the fish consumed less quantity of diet 6 due to easy distrigration and sinking tendency of the feed sample. But the little quantity of diet 6 consumed was well utilized.




Fig. 1a. Weekly feeds sampled consumed




Fig. 1b. Pattern of diff. feeds consumed and weight gained




Fig. 1c. Level of sampled feeds consumed




Fig. 2a. Pattern of feed conversion ratio with diff. feeds sampled




Fig. 2b. Feed conversion ratio of sampled fish feeds



Fig. 3. Length-weight relationships




Fig. 4. Cost per tonne of different feeds

Fig. 3 shows the length -weight relationship of the test organism in the experiment. It was obvious tha fish had relatibvely uniform and steady lenth-weight relationship in the first two weeks. After which there were sharp drops in the growth perttern of fish in floating types. The sinking type progressively sustained the positive allometry growth better than floating feeds. This may be attributed to the percentage crude protein content of each feed relative to the size of fish.

The analysis was also carried out on the cost per tonne of different feeds sampled. Fig. 4 showed that diet 6 had the least cost per tonne while diet 5 had the highest cost. Relating the cost of individual diet with feed convesion ratio, it was discovered that diet 6 had the least cost and highest feed conversion ratio while diet 5 had the highest cost with least feed conversion ratio. It could be deduced that diet 6 is most economical diet than other extruded floating types with concomitant marginal profits.

4. CONCLUSION

The results of this study had shown that there are falsifications in crude protein percentages. The actual crude protein percentage in each feed was established through proximate analysis, response of fish growth and feed utilization efficiency. Generally, floating feeds performed excellently well in terms of weight gained and length-weight relationships for the first two weeks as evidenced in the experiment. However, the body weight gained dropped sharply and did not commensurate with the total feeds consumed

over time. The control diet (DT₆) was least consumed with steady body weight gained and positive weight length-weight relationship and had best feed conversion ratio.

5. RECOMMENDATION

According to the experiment conducted, it revealed that floating feeds specifically (DT1) and (DT2) are good and preferable to be used in early 2-4 weeks after which compounded feed is recommended for least feed consumption, steady body weight-gained and for high cost effectiveness.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tsevis AA, Azzaydi TA. Effect of feeding regime on selected species of fish Article Publicaiton of FISON. 20000 Agora site; 2000.
2. Jamiu DM, Ayinla OA. Potential for the development of aquaculture in Africa. NAGA 2003;26(3):6-13.
3. Akinrotimi. Locally produced fish feed, potentials for aquaculture development in sub-sharan. African Journal of Agricultural Resources. 2007;297.
4. Olakunle O. Homestead pond management, Department of wild life and fisheries management University of Ibadan Oyo State, Nigeria; 2000.

5. Alatise SP, Adedokun MA, Adedokun OB, Ajiboye GE. Effects of boiled jatropha kernel meal as a substitute for soyabeans meal in diet of African Mud Catfish (*Clarias gariepinus*). Journal of Fisheries and Aquatic Scien; 2014. [ISSN 1816-4927] DOI: 10, 3923/jfas.
6. Anetekhai MA, Akin-Oriola GA, Aderinola OJ, Akintola S.L. Steps ahead for aquaculture development in sub-saharan Africa-the case of Nigeria. Aquaculture. 2004;239:237-248.
7. Hect T. Review of feed and fertilizer for sustainable Aquaculture in Sub-saharan Africa. In MR Hasan, T Hecht, SS De Silva, AGJ Tacon (eds). Study and analysis of feed and fertilizer for sustainable Aquaculture development. FAO Fisheries Technical Paper No/ 497 Rome FAO. 2007;77-109.
8. Ayinla OA. Analysis of feeds and fertilizer for sustainable aquaculture development in Nigeria. In MR Hassan, T Hecht, SS De Silua, AGJ Tacon (eds). Study and analysis of feed and fertilizer for sustainable Aquaculture development. FAO Fisheries Technical paper No. 497. Rome. FAO. 2007;453-470.
9. Ayinla OA, Akande GR. Growth response of *Clarias gariepinus* (Burchell, 1822) on silage base diets.NIOMR. Technical Paper. 1988;37:19.
10. Achionye – Nzeh CG, Qgidiolu O, Salmi S. Growth response of juveniles of *Clarias anguila* diet formulated, Ciriunaforda within laboratory. Nigerian Journal of Pure and Applied Sciences. 2002;17:1253-1256.
11. Fagbenro OA, Arowosoge IA. Replacement value of some household wastes as energy substitute in low rearing catfish, in south-western Nigeria. BioresourceTechnol. 2002;37:197-203.
12. Otubusin SO, Ogunleye FO, Agbebi OI. Feeding trials using local protein sources to replace fish meal in pelleted feed in catfish *Clarias gariepinus* (Burchell, 1822) culture. European Journal of Scientific Research. 2009;31(1):142-174.
13. Amisah S. Oteng MA, Ofori JK. Growth performance of the African Catfish *Clarias gariepinus* Fed varying inclusion level of *Leavcaena leaucocephala* leaf meal. Journal of Applied Science and Environmental Management. 2009;13(1):21-26.
14. Sotolu AA. Comparative utilization of fish waste meal with imported fish meal by Africa cat fish (*Clarias gariepinus*). American Europeans Journal of Scientific Research. 2009;4(4):225-289.
15. Sotolu AO. Feed utilization and biochemical characteristic of *Clariasgariepinus* (Qurchz//, 1% 22) finger ling fed diet containing fish oil and vegetable oil as a total replacement. World Journal of Fish and Marine Science. 2010;2(2):93-98.
16. SPSS. Software Program of Statistical Analysis. Version 8.0, SPSS Inc., Chicago, IL., USA; 1999.
17. APHA/AWWAWPCF. Standard methods for the examination of water and wastewaters. The 20th Edition. American Public Health Association American Water works. Association and water Pollution Control Federation; Washington Inc. 1999;460-472.
18. Adedokun MA, Tairu HM, Adeosun O, Ajibola O. Assessment of the optimal replacement Levels of maize with water lettuce leaf (*Pistia stratiotes*) based diets for *Clarias gariepinus*. Journal of Fisheries Sciences. Com. 2017;11(2):028-035.
19. Ajani EK, Akinwale OA. Recommended water quality for warm water Fishes in Fish Farming. University of Ibadan. Agric Res. 2001;3:19-23.

© 2019 Adedokun et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50215>